

PD-L1
Recombinant Rabbit Monoclonal Antibody
Product Datasheet

Catalog# BX00006

Clone# RR604

Predicted Molecular Wt: 33kDa

Purity: ProA affinity purified IgG

Species Cross-reactivity: Human

Form: Liquid


Species cross-reactivity determined by WB

Swissprot ID: Q9NZQ7

Applications: WB IHC-P IF/ICC FC IP

Background:

Involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PDCD1-independent manner. Interaction with PDCD1 inhibits T-cell proliferation and cytokine production.



All lanes: Anti-PD-L1 antibody at 1:5,000 dilution

Predicted MW: 33 kDa

Observed MW: 40-50 kDa

Lane 1: HEK293 Overexpression of HuPD-L1

Lane 2: HEK293

Lysates at 5 µg per lane

2nd Ab:

GAR HRP(H+L) 1:10,000

Exposure: 100s

Immunogen:

Recombinant full length protein corresponding to Human PD-L1. The immunogen contains the specific extracellular domain of huPD-L1 (F19-T239).

Storage Buffer:

PBS 59%, Sodium azide 0.01%, Glycerol 40%, BSA 0.05%.

Storage conditions:

-20°C.

Storage instructions:


Shipped on blue ice. Upon delivery, aliquot, and store at -20°C. Avoid freeze / thaw cycles.

Recommended Dilutions:


WB: 1:2,500 - 1:5,000
 IHC-P: 1:100 - 1:200
 IF/ICC: 1:2,000 - 1:10,000
 FC: 1:50 - 1:200
 IP: 1:30

Background References:

- Adam J., Ann Pathol. 2016 Jan;36(1):94-102.
- Lim SH., Expert Opin Biol Ther. 2016;16(3):397-406.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human lung adenocarcinoma tissue labelling PD-L1 with RR604 at 1:200. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9.0.




RR604 staining 293 cells transfected with PD-L1 gene by IF/ICC (immunofluorescence/immunocytochemistry). Cells were fixed with paraformaldehyde, permeabilized with 0.1% Triton X-100 and blocked with 10% goat serum for half an hour at room temperature. Samples were incubated with primary antibody (1:10,000) at 4°C. An Alexa Fluor® 488-conjugated Goat Anti-Rabbit IgG polyclonal was used as the secondary antibody (1:500). DAPI (blue) was used as the nuclear counter stain.

Control: PBS and secondary antibody, An Alexa Fluor® 488-conjugated Goat Anti-Rabbit IgG (1:500).



Overlay histogram showing 293 cells transfected with PD-L1 gene stained with RR604. The cells were then incubated in the antibody (RR604, 1:200 dilution) in 1x PBS/1% BSA for 30 min at room temperature. The secondary antibody used was a Goat Anti-Rabbit Alexa Fluor® 488 (IgG H+L) at 1:2,000 dilution for 20 min at room temperature. Unlabelled sample (Black) was used as a control.



PD-L1 was immunoprecipitated from 0.2mg of 293 whole cells lysate transfected with PD-L1 gene with RR604 at 1:30 dilution.

2nd Ab: GAR HRP for IP 1:500

Lane 1: RR604 IP in 293 whole cell lysate transfected with PD-L1 gene

Lane 2: PBS instead of RR604 in 293 whole cell lysate transfected with PD-L1 gene

Lane 3: 293 whole cell lysate transfected with PD-L1 gene, 2 µg (input)

Exposure: 10s

Product QC'd by:



For research use only. Not for use in diagnostic or therapeutic applications.