

TROP-2 Recombinant Rabbit Monoclonal Antibody Product Datasheet

Catalog# BX50231

Clone# BP6208

Predicted Molecular Wt: 36kDa
Species Cross-reactivity: Human
Applications: IHC-P

Purity: ProA affinity purified IgG
Form: Liquid
Swissprot ID: P09758

Background:

Trop-2 or Tumor-associated calcium signal transducer 2, also known as epithelial glycoprotein-1 antigen (EGP-1), is a protein that in humans is encoded by the TACSTD2 gene. This transmembrane glycoprotein functions in a variety of cell signaling pathways and was first elucidated as a transducer of an intracellular calcium signal. Trop-2 expression has been demonstrated to depend on a large variety of transcription factors. Trop-2 is involved in several cell signaling pathways, of which many are associated with tumorigenesis.

Subcellular location:

Membrane

Recommended method:

Heat induced epitope retrieval with Tris-EDTA buffer (pH 9.0), primary antibody incubate at RT (18°C-25°C) for 30 minutes.

Immunogen:

Synthetic peptide. This information is proprietary to Biolynx.

Storage Buffer:

PBS 59%, Sodium azide 0.01%, Glycerol 40%, BSA 0.05%.

Storage conditions:

-25°C to -18°C

Storage instructions:


Shipped on blue ice. Upon delivery, aliquot, and store at -25°C to -18°C. Avoid freeze / thaw cycles.

Recommended Dilutions:

IHC-P: 1:100-1:200

Background References:

1. McDougall, A.R. et al. (2015) Dev Dyn 244, 99-109.
2. Shvartsur, A. and Bonavida, B. (2015) Genes Cancer 6, 84-105.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human esophageal carcinoma tissue labelling TROP-2 with BP6208. Heat mediated antigen retrieval was performed using Tris/EDTA buffer pH 9.0

Product QC'd by:



For research use only. Not for use in diagnostic or therapeutic applications.